


N-Benzotriazol-1-yl-methyl-1,2,3,4-tetrahydroisoquinoline

Cornelia Locher*, Sara Baldey and Naseem Peerzada*

Faculty of Science, Northern Territory University, Darwin, N.T. 0909, Australia - ph: 0061-8-8946 6850, fax: 0061-8-8946 6847, e-mails (Cornelia Locher) clocher@banda.ntu.edu.au, (Sara Baldey) sbaldey@banda.ntu.edu.au, (Naseem Peerzada) peerzadan@banks.ntu.edu.au

Received: 16 October 1998 / Published: 22 February 1999



Benzotriazole-mediated heteroalkylations have been explored for many synthetic pathways. The principle of an aminoalkylation (or Mannich reaction) with primary or secondary amines was applied to the reaction of 1-hydroxymethylbenzotriazole [1] with 1,2,3,4-tetrahydroisoquinoline [2] which yielded very high quantities of crystalline *N*-benzotriazol-1-ylmethyl-1,2,3,4-tetrahydroisoquinoline [3].

1,2,3,4-Tetrahydroisoquinoline (19.9mmol, 2.64g) was added to 1-hydroxymethylbenzotriazole (2.96g, 19.9mmol) with stirring. Ethanol (5mL) was added and the mixture heated. The reaction was very fast and initial crystallisation of the product started already after a few seconds. More ethanol (15mL) was added and the solution refluxed with stirring for 30min. The mixture was kept at -5 °C over night, the final product (5.14g, 98%) was collected under reduced pressure and washed with cold ethanol. It was recrystallised from ethanol and yielded *N*-benzotriazol-1-ylmethyl-1,2,3,4-tetrahydroisoquinoline [3] as white needles.

M.p. 155-157 °C (EtOH, uncorrected).

UV $I_{\max}(\text{nm}; \text{EtOH}) / \epsilon (\text{dm}^3 \text{mol}^{-1} \text{cm}^{-1})$ 206 / 27060, 253 / 6970 and 273 / 5808.

IR $\nu_{\max}(\text{cm}^{-1}; \text{Nujol})$ 1260, 1210, 1150, 1090, 1050, 955, 940, 740.

$^1\text{H-NMR}$ d_{H} (200 MHz; CDCl_3 ; Me_4Si) 8.08 (1H, d, J 4), 7.68 (1H, d, J 4), 7.52 (1H, t, J 7), 7.38 (1H, t, J 7), 7.10-6.90 (4H, m), 5.64 (2H, s, NCH_2N), 3.86 (2H, s, ArCH_2N), 2.96-2.91 (4H, m, $\text{ArCH}_2\text{CH}_2\text{N}$).

$^{13}\text{C-NMR}$ d_{C} (50 MHz; CDCl_3) 29.0 ($\text{ArCH}_2\text{CH}_2\text{N}$), 48.4 (ArCH_2N), 52.3 ($\text{ArCH}_2\text{CH}_2\text{N}$), 69.1 (NCH_2N), 110.1, 118.4, 120.1, 124.1, 125.9, 126.4, 126.6, 127.7, 128.9, 133.6, 133.7, 134.0, 146.1.

Analysis Found ($\text{C}_{16}\text{H}_{16}\text{N}_4$, 264.33, Calc.) C 72.6 (72.7), H 6.2 (6.1), N 21.2 (21.2).

Acknowledgments: One of the authors (C.L.) would like to thank the Northern Territory University for the receipt of a University Postgraduate Research Scholarship.

References

1. Katritzky, A.R.; Lan, X.; Yang, J.Z.; Denisko, O.V. Properties and Synthetic Utility of

N-Substituted Benzotriazoles. *Chem. Rev.* **1998**, *98*, 409.

2. Katritzky, A.R.; Rachwal, S.; Rachwal, B. The Chemistry of Benzotriazole. Part 3. The Aminoalkylation of Benzotriazole. *J. Chem. Soc., Perkin Trans. I* **1987**, 799.

3. Viktor, M.; Rudolf, K. 1-Benzimidazolyl-1-benztriazolylmethane. *Molecules* **1997**, *2*, M12.

Sample Availability: Available from the authors (C.L. or N.P.) and from MDPI. [MDPI Reg. No. 15927](#).

©1999 MDPI. All rights reserved. *Molecules* website <http://www.mdpi.org/molecules/>